

June 4, 1953

20862

SUBJECT: New Products - May 1953

Domestic

Aureomycin Triple Sulfas

Description: Each tablet contains:
Aureomycin HCL 125 mg.
Sulfadiazine 167 mg.
Sulfamerazine 167 mg.
Sulfamethazine 167 mg.

Indications: For the treatment of shigella dysentery where the action of Aureomycin may be insufficient to combat the organism and for the treatment of gonorrhoea.

Administration: For shigella dysentery: average daily adult dose is 2 tablets four times daily
For gonorrhoea: recommended dose is 2 tablets followed by 1 tablet at 6 hour intervals for 2 doses. This course may be repeated if necessary.

Form Supplied: 25X1X6 Bottles of 12's

Source: [REDACTED]

Depo-Cer-O-Cillin

Description: A water-insoluble, crystalline salt of penicillin O (allylmercapto-methyl penicillin). The addition of 4.2 cc. of sterile water for injection or sterile isotonic solution of sodium chloride provides 5 cc. of a suspension containing 300,000 units of Depo-Cer-O-Cillin per cc. The dry form is stable for 36 months at room temperature. Suspensions are stable at room temperature for one week and at refrigerator temperature for three weeks.

Indications: For the treatment of acute staphylococccic, streptococccic, and pneumococccic infections; bacteremia and gonorrhoea.

Administration: Average dose is 300,000 units by intramuscular injection, or as directed by physician.

Form Supplied: 25X1X6 Sterile vials, containing 1,500,000 units.

Source: [REDACTED]

Liquocillin

Description: Each fluid drachm contains: 100,000 units potassium penicillin G.

Indications: Administering penicillin by mouth to infants and young children.


Form Supplier: Bottles of 2 fluid ozs.

Source: [REDACTED]

25X1X6

E.R. Squibb & Sons, New York has changed the name of Penfonylin to Pentid-Sulfas. They contain: Squibb 200,000 Units penicillin G potassium with 0.5 Gm. Meth-Dia-Mer Sulfonamide Tablets.

25X1X6



al.

* Competitive with Pfizer's Permapen

Crystamycin

Description:

Contains:

crystalline penicillin G with a mixture of streptomycin and dihydrostreptomycin sulfates.

Indications:

Used for treating infections due to organisms relatively resistant to the antibiotics used singly. Its use is indicated where the identity and degree of sensitivity of the infecting strains have not been determined, and a mixed infection or the presence of a resistant strain is suspected (as possibly in infections of the urinary tract, in peritonitis and in infected wounds).

25X1X6

Form Supplied:

Box of ten single dose vials

Source:

Diazil-Penicillina

Description:

Each tablet contains:

diazil (p. aminobenzosulphanilamino)	4.60
dimethylpyrimidine	0.50 Gm.
penicillin	100,000 Units

Indications:

Pneumonia, bronchial pneumonia etc.

Administration:

2-3 tablets, 3 to 4 times a day

Form Supplied:

Tubes of 12 tablets

25X1X6

* Dimycin

Description:

Dry mixture of streptomycin sulfate and dihydrostreptomycin sulfate in equal parts by potency.

Indications:

Used in the treatment of some forms of tuberculosis, mainly in cases where the disease is progressive, of relatively short duration, also in certain non-tubercular infections not amenable to penicillin.

25X1X6

Form Supplied:

Box of ten 1 gram vials and 5 gram vials

Pansulfa Penicillina

Description:

Each tablet contains:

Sulphadiazine	0.166 gms.
Sulphamerazine	0.166 gms.
Sulphathiazole	0.166 gms.
Penicillin G	150,000 Units

Indications:

All coccus infections

Administration:

2 tablets, 3-4 times a day

Form Supplied:

Tablets and suppositories

* Competitive with Pfizer's Combistrep

Rettocillina

Description: Each suppository contains:
Aureomycin 0.125 Gm. or .250 Gm.
25X1X6 Penicillin 12,000 Units
Indications: All the infections caused by germs susceptible to Aureomycin.
Administration: 3-6 suppositories a day
Form Supplied: Boxes of 3's

25X1X6

Sintomicetina Ointment (Chloramphenicol)

Description: Ointment - 5%
Indications: Skin infections, acne, anthrax, eczema, herpes, warts
Form Supplied: Container of 5 Gm. ointment

Sintomicetina Ophthalmic Solution (Chloramphenicol)

25X1X6

Indications: Acute conjunctivitis, catarrhal conjunctivitis (Morax-Axenfeld, Koth-Weeks), gonococcal conjunctivitis, purulent and pseudomembranous conjunctivitis caused by staphylococci, pneumococci or streptococci.
Administration: A few drops 3 or 5 times a day
Form Supplied: Vials of 20 cc. with special pipette

Streptosuccinil

25X1X6

Description: Streptomycin and Succinyl-Sulfatiazol in apple pectin
Indications: Intestinal infections in infants

Veterinary

Vetstrep

25X1X6

Description: Oral veterinary streptomycin sulfate
Indications: Treatment of swine enteritis, calf scours and other enteritis
Administration: Orally, grains of Vetstrep may be given to animals in drinking water or milk
Form Supplied: Bottles of 15, 75 and 750 Gms.
Source: Merck & Co., Rahway, N. J.

lete line

of food substances. They have quoted the following prices:

Distafed riboflavin supplement		
5,000 microgm. per gm.	1 lb.	\$.65
	10 lb.	5.83
	50 lb.	28.00
1,000 microgm. per gm.	1 lb.	.26
	10 lb.	1.87
	50 lb.	8.17

Distafed vitamin B12 supplement		
15 microgm. per gm.	1 lb.	.84
	10 lb.	7.70

Distafeed antibiotic supplement

1 gm. procaine penicillin per lb.

1 lb. \$.44

10 lb. 3.73

subject to discount of 25 per cent to retail pharmacists
and available only through wholesalers.